ELSEVIER

Molecular Cell Research

Vol. 1356, 1997

BIOCHIMICA ET BIOPHYSICA ACTA
BBA

Cumulative Contents

Regular Section

Information for Contributors *vii*

Regular papers

Porcine $\alpha_{1,3}$-galactosyltransferase: tissue-specific and regulated expression of splicing isoforms
B. Vanhove, F. Goret, J.-P. Soulillou and C. Pourcel (France) 1

Loading with oxidised low density lipoprotein alters endocytic and secretory activities of murine macrophages
E.J. Bolton, W. Jessup, K.K. Stanley and R.T. Dean (Australia) 12

An amino terminal prosequence is required for efficient synthesis of *S. cerevisiae* **a**-factor
G.E. Quinby and R.J. Deschenes (USA) 23

The inhibitory effect of vitamin E on pulmonary polyamine biosynthesis, cell proliferation and carcinogenesis in mice
Y. Yano, T. Yano, M. Uchida, A. Murakami, M. Ogita, T. Ichikawa, S. Otani and K. Hagiwara (Japan) 35

Differential induction of constitutive and inducible nitric oxide synthases by distinct inflammatory stimuli in bovine aortic endothelial cells
Y. Kaku, H. Nanri, T. Sakimura, K. Ejima, A. Kuroiwa and M. Ikeda (Japan) 43

Analysis of expression of an alternative La (SS-B) cDNA and localization of the encoded N- and C-terminal peptides
M. Bachmann, D. Grölz, H. Bartsch, R.R. Klein and H. Tröster (Germany) 53

Protein kinase A-dependent activation of PDE4 (cAMP-specific cyclic nucleotide phosphodiesterase) in cultured bovine vascular smooth muscle cells
D. Ekholm, P. Belfrage, V. Manganiello and E. Degerman (Sweden, USA) 64

Ca^{2+}-dependent membrane bound protein fraction from rabbit gastric mucosa contains a protein whose histidyl residue is phosphorylated
T. Urushidani and T. Nagao (Japan) 71

Subsection

Regular papers

Methylglyoxal-modified arginine residues — a signal for receptor-mediated endocytosis and degradation of proteins by monocytic THP-1 cells
M.E. Westwood, O.K. Argirov, E.A. Abordo and P.J. Thornalley (UK) 84

Immunocytochemical detection of estrogen receptors in bone cells using flow cytometry
T. Štulc, D. Klement, J. Kvasnička and J.J. Štěpán (Czech Republic) 95

A novel inhibitory action of wheat germ agglutinin on phospholipase C in HEL and MEG-01 cell lines
E. Rezaei, L. Quek and S.P. Watson (UK) 101

Enhancement of tissue-type plasminogen activator (t-PA) activity by purified t-PA receptor expressed in human endothelial cells
H. Fukao, S. Ueshima, T. Takaishi, K. Okada and O. Matsuo (Japan) 111

Information for Contributors *vii*

Regular Section

Regular papers

Inhibitors of actin polymerization and calmodulin binding enhance protein kinase C-induced translocation of MARCKS in C6 glioma cells
D.N. Douglas, H.-S. Fink, S.D. Rosé, N.D. Ridgway, H.W. Cook and D.M. Byers (Canada) 121

Nitric oxide synthase and arginase in cells isolated from the rat gastric mucosa
C.R. Byrne, K.J. Price, J.M. Williams, J.F. Brown, P.J. Hanson and B.J.R. Whittle (UK) 131

Electrogenic ion transport in the mouse endometrium: functional aspects of the cultured epithelium
H.C. Chan, C.Q. Liu, S.K. Fong, S.H. Law, P.S. Leung, P.Y. Leung, W.O. Fu, S.B. Cheng Chew and P.Y.D. Wong (Hong Kong, China) 140

Differential expression of proteins regulating cell cycle progression in growth vs. differentiation
Z.-r. Li, R. Hromchak and A. Bloch (USA) 149

Role of prostaglandin E_2 in regulation of low and high water osmotic permeability in frog urinary bladder
R.G. Parnova, E.I. Schakhmatova, S.A. Plesneva, E.V. Getmanova, E.V. Korolev, Y.Y. Komissarchik and Yu.V. Natochin (Russia) 160

Overproduction of urokinase-type plasminogen activator is regulated by phospholipase D- and protein kinase C-dependent pathways in murine mammary adenocarcinoma cells
J.A. Aguirre Ghiso, D.F. Alonso, E.F. Farías and E. Bal de Kier Joffé (Argentina) 171

Cationic liposome-mediated expression of HIV-regulated luciferase and diphtheria toxin a genes in HeLa cells infected with or expressing HIV
K. Konopka, G.S. Harrison, P.L. Felgner and N. Düzgüneş (USA) 185

Effect of caffeine on mucus secretion and agonist-dependent Ca^{2+} mobilization in human gastric mucus secreting cells
E. Hamada, T. Nakajima, Y. Hata, H. Hazama, K. Iwasawa, M. Takahashi, S.-i. Ota and M. Omata (Japan) 198

Subsection

Regular papers

A characterization of permolybdate and its effect on cellular tyrosine phosphorylation, gap junctional intercellular communication and phosphorylation status of the gap junction protein, connexin43
S.-O. Mikalsen and O. Kaalhus (Norway) 207

Stimulation of atrial natriuretic peptide receptor/guanylyl cyclase - A signaling pathway antagonizes the activation of protein kinase C-α in murine Leydig cells
R. Kumar, T.W. Von Geldern, R.A. Calle and K.N. Pandey (USA) 221

Expression and signal transduction of the glucagon receptor in βTC3 cells
J. Li, J.N. Larocca, A.G. Rodriguez-Gabin and M.J. Charron (USA) 229

T cell antigen receptor dependent signalling in human lymphocytes: cholera toxin inhibits interleukin-2 receptor expression but not interleukin-2 synthesis by preventing activation of a protein kinase C isotype, PKC-α
M. Szamel, U. Ebel, P. Uciechowski, V. Kaever and K. Resch (Germany) 237

Information for Contributors *vii*

Regular Section

Short sequence-papers

Presence of multiple functional polyadenylation signals in the 3'-untranslated region of human corticotropin receptor cDNA
A. Penhoat, D. Naville, C. Jaillard, P. Durand and M. Bégeot (France) 249

Isolation of a *Xenopus laevis* genomic clone representing a novel *N*-cadherin related gene
S.A. Burbidge, P.M. Webber and E.A. Jones (UK) 253

Tyrosine phosphorylation-and epidermal growth factor-dependent regulation of the sodium-coupled amino acid transporter B^0 in the human placental choriocarcinoma cell line JAR
V. Torres-Zamorano, R. Kekuda, F.H. Leibach and V. Ganapathy (USA) 258

Suppression of basement membrane type IV collagen degradation and cell invasion in human melanoma cells expressing an antisense RNA for MMP-1
M. Durko, R. Navab, H.R. Shibata and P. Brodt (Canada) 271

Decreased transcript expression coincident with impaired glycosylation in the β_2-adrenergic receptor gene does not result from differences in the primary sequence
R.J. Hughes, M. Pasillas, J. Saiz, J. Jasper and P.A. Insel (USA, Spain) 281

Catecholamines are required for androgen-induced ODC expression but not for hypertrophy of mouse kidney
M. Manteuffel-Cymborowska, M. Pęska, W. Chmurzyńska and B. Grzelakowska-Sztabert (Poland) 292

Subsection

Regular papers

Cloning and functional analysis of a polymorphic variant of the ovine Mel 1a melatonin receptor
P. Barrett, S. Conway, R. Jockers, A.D. Strosberg, B. Guardiola-Lemaitre, P. Delagrange and P.J. Morgan (UK, France) 299

Dissociation of tyrosine phosphorylation and activation of phosphoinositide phospholipase C induced by the protein kinase C inhibitor Ro-31-8220 in Swiss 3T3 cells treated with platelet-derived growth factor
E.-J. Yeo, J.J. Provost and J.H. Exton (USA) 308

The thrombin receptor in adrenal medullary microvascular endothelial cells is negatively coupled to adenylyl cyclase through a G_i protein
V.G. Manolopoulos, J.W. Fenton II and P.I. Lelkes (USA) 321

Inhibition of protein tyrosine kinases or protein kinase C prevents nonspecific killer T lymphocyte-mediated tumoricidal activity
B.H. Stewart and D.W. Hoskin (Canada) 333

The gastrin-releasing peptide receptor is differentially coupled to adenylate cyclase and phospholipase C in different tissues
L.J. García, T.K. Pradhan, H.C. Weber, T.W. Moody and R.T. Jensen (Spain, USA) 343

Cumulative Contents, Vol. 1356 355

Author Index 358